

OXFORD CAMBRIDGE AND RSA EXAMINATIONS
Sixth Term Examination Papers
administered on behalf of the Cambridge Colleges

MATHEMATICS I

9465

Friday

29 JUNE 2001

Afternoon

3 hours

Additional materials:

Answer paper

Graph paper

Formulae booklet

Candidates may not use electronic calculators

TIME 3 hours

INSTRUCTIONS TO CANDIDATES

Write your name, Centre number and candidate number in the spaces provided on the answer paper/answer booklet.

Begin each answer on a new page.

INFORMATION FOR CANDIDATES

All questions carry equal marks.

Extra credit is given for substantially complete answers.

You will be assessed on the six questions for which you gain the highest marks. There is no restriction of choice.

You are provided with Mathematical Formulae and Tables.

The use of electronic calculators is not permitted.

This question paper consists of 6 printed pages and 2 blank pages.

Section A: Pure Mathematics

- 1 The points A , B and C lie on the sides of a square of side 1 cm and no two points lie on the same side. Show that the length of at least one side of the triangle ABC must be less than or equal to $(\sqrt{6} - \sqrt{2})$ cm.

- 2 Solve the inequalities

(i) $1 + 2x - x^2 > 2/x \quad (x \neq 0),$

(ii) $\sqrt{(3x + 10)} > 2 + \sqrt{(x + 4)} \quad (x \geq -10/3).$

- 3 Sketch, without calculating the stationary points, the graph of the function $f(x)$ given by

$$f(x) = (x - p)(x - q)(x - r),$$

where $p < q < r$. By considering the quadratic equation $f'(x) = 0$, or otherwise, show that

$$(p + q + r)^2 > 3(qr + rp + pq).$$

By considering $(x^2 + gx + h)(x - k)$, or otherwise, show that $g^2 > 4h$ is a sufficient condition but not a necessary condition for the inequality

$$(g - k)^2 > 3(h - gk)$$

to hold.

- 4 Show that $\tan 3\theta = \frac{3 \tan \theta - \tan^3 \theta}{1 - 3 \tan^2 \theta}.$

Given that $\theta = \cos^{-1}(2/\sqrt{5})$ and $0 < \theta < \pi/2$, show that $\tan 3\theta = 11/2.$ ✓

Hence, or otherwise, find all solutions of the equations

(i) $\tan(3 \cos^{-1} x) = 11/2,$

(ii) $\cos(\frac{1}{3} \tan^{-1} y) = 2/\sqrt{5}.$

5 Show that (for $t > 0$)

$$(i) \quad \int_0^1 \frac{1}{(1+tx)^2} dx = \frac{1}{(1+t)},$$

$$(ii) \quad \int_0^1 \frac{-2x}{(1+tx)^3} dx = -\frac{1}{(1+t)^2}.$$

Noting that the right hand side of (ii) is the derivative of the right hand side of (i), conjecture the value of

$$\int_0^1 \frac{6x^2}{(1+x)^4} dx.$$

(You need not verify your conjecture.)

6 A spherical loaf of bread is cut into parallel slices of equal thickness. Show that, after any number of the slices have been eaten, the area of crust remaining is proportional to the number of slices remaining.

A European ruling decrees that a parallel-sliced spherical loaf can only be referred to as 'crusty' if the ratio of volume V (in cubic metres) of bread remaining to area A (in square metres) of crust remaining after any number of slices have been eaten satisfies $V/A < 1$. Show that the radius of a parallel-sliced spherical loaf must be less than $2\frac{2}{3}$ metres.

[The area A and volume V formed by rotating a curve in the x - y plane round the x -axis from $x = -a$ to $x = -a + t$ are given by

$$A = 2\pi \int_{-a}^{-a+t} y \left(1 + \left(\frac{dy}{dx} \right)^2 \right)^{\frac{1}{2}} dx, \quad V = \pi \int_{-a}^{-a+t} y^2 dx.]$$

- 7 In a cosmological model, the radius R of the universe is a function of the age t of the universe. The function R satisfies the three conditions:

$$R(0) = 0, \quad R'(t) > 0 \text{ for } t > 0, \quad R''(t) < 0 \text{ for } t > 0, \quad (*)$$

where R'' denotes the second derivative of R . The function H is defined by

$$H(t) = \frac{R'(t)}{R(t)}.$$

- (i) Sketch a graph of $R(t)$. By considering a tangent to the graph, show that $t < 1/H(t)$.
 - (ii) Observations reveal that $H(t) = a/t$, where a is constant. Derive an expression for $R(t)$. What range of values of a is consistent with the three conditions $(*)$?
 - (iii) Suppose, instead, that observations reveal that $H(t) = bt^{-2}$, where b is constant. Show that this is not consistent with conditions $(*)$ for any value of b .
- 8 Given that $y = x$ and $y = 1 - x^2$ satisfy the differential equation

$$\frac{d^2y}{dx^2} + p(x) \frac{dy}{dx} + q(x)y = 0, \quad (*)$$

show that $p(x) = -2x(1 + x^2)^{-1}$ and $q(x) = 2(1 + x^2)^{-1}$.

Show also that $ax + b(1 - x^2)$ satisfies the differential equation for any constants a and b .

Given instead that $y = \cos^2(x^2/2)$ and $y = \sin^2(x^2/2)$ satisfy the equation $(*)$, find $p(x)$ and $q(x)$.

Section B: Mechanics

- 9 A ship sails at 20 kilometres/hour in a straight line which is, at its closest, 1 kilometre from a port. A tug-boat with maximum speed 12 kilometres/hour leaves the port and intercepts the ship, leaving the port at the latest possible time for which the interception is still possible. How far does the tug-boat travel?

- 10 A gun is sited on a horizontal plain and can fire shells in any direction and at any elevation at speed v . The gun is a distance d from straight railway line which crosses the plain, where $v^2 > gd$. The gunner aims to hit the line, choosing the direction and elevation so as to maximize the time of flight of the shell. Show that the time of flight, T , of the shell satisfies

$$g^2 T^2 = 2v^2 + 2(v^4 - g^2 d^2)^{\frac{1}{2}}.$$

- 11 A smooth cylinder with circular cross-section of radius a is held with its axis horizontal. A light elastic band of unstretched length $2\pi a$ and modulus of elasticity λ is wrapped round the circumference of the cylinder, so that it forms a circle in a plane perpendicular to the axis of the cylinder. A particle of mass m is then attached to the rubber band at its lowest point and released from rest.

- (i) Given that the particle falls to a distance $2a$ below the axis of the cylinder, but no further, show that

$$\lambda = \frac{9\pi mg}{(3\sqrt{3} - \pi)^2}.$$

- (ii) Given instead that the particle reaches its maximum speed at a distance $2a$ below the axis of the cylinder, find a similar expression for λ .

Section B: Probability and Statistics

- 12 Four students, Arthur, Bertha, Chandra and Delilah, exchange gossip. When Arthur hears a rumour, he tells it to one of the other three without saying who told it to him. He decides whom to tell by choosing at random amongst the other three, omitting the ones that he knows have already heard the rumour. When Bertha, Chandra or Delilah hear a rumour, they behave in exactly the same way (even if they have already heard it themselves). The rumour stops being passed round when it is heard by a student who knows that the other three already have already heard it.

Arthur starts a rumour and tells it to Chandra. By means of a tree diagram, or otherwise, show that the probability that Arthur rehears it is $3/4$.

Find also the probability that Bertha hears it twice and the probability that Chandra hears it twice.

- 13 Four students, one of whom is a mathematician, take turns at washing up over a long period of time. The number of plates broken by any student in this time obeys a Poisson distribution, the probability of any given student breaking n plates being $e^{-\lambda}\lambda^n/n!$ for some fixed constant λ , independent of the number of breakages by other students. Given that five plates are broken, find the probability that three or more were broken by the mathematician.
- 14 On the basis of an interview, the N candidates for admission to a college are ranked in order according to their mathematical potential. The candidates are interviewed in random order (that is, each possible order is equally likely).
- (i) Find the probability that the best amongst the first n candidates interviewed is the best overall.
 - (ii) Find the probability that the best amongst the first n candidates interviewed is the best or second best overall.

Verify your answers for the case $N = 4$, $n = 2$ by listing the possibilities.